

Liver cancer

1993-2022

(ICD10 codes: C22)



Northern Ireland
Cancer Registry

Northern Ireland Cancer Registry, 2025

An official statistics publication

ABOUT THIS REPORT

Contents

This report includes information on incidence of liver cancer as recorded by the Northern Ireland Cancer Registry (NICR). Incidence data is available annually from 1993 to 2022, however in order to provide stable and robust figures the majority of information presented in this report is based upon the average number of cases diagnosed in the last five years.

Methodology

The methodology used in producing the statistics presented in this report, including details of data sources, classifications and coding are available in the accompanying methodology report available at: www.qub.ac.uk/research-centres/nicr/CancerInformation/official-statistics.

Official statistics

The incidence, prevalence and survival statistics in this publication are designated as official statistics signifying that they comply with the Code of Practice for Official Statistics. Further information on this code is available at code.statisticsauthority.gov.uk.

Cancer mortality data

The NI Statistics and Research Agency (NISRA) is the official statistics provider of cancer mortality data in Northern Ireland. However, for completeness, data on cancer mortality is also provided in this report. While analysis is conducted by NICR staff, the original data is provided courtesy of the General Register Office (NI) via the Department of Health.

Reuse of information

The information in this report (and any supplementary material) is available for reuse free of charge and without the need to contact NICR. However, we request that NICR is acknowledged as the source of any reused information. The following reference is recommended:

Northern Ireland Cancer Registry 2025. Liver cancer: 1993-2022. Available at: www.qub.ac.uk/research-centres/nicr

Further information

Further information is available at: www.qub.ac.uk/research-centres/nicr

Phone: +44 (0)28 9097 6028 **e-mail:** nicr@qub.ac.uk

Acknowledgements

The Northern Ireland Cancer Registry (NICR) uses data provided by patients and collected by the health service as part of their care and support.


NICR is funded by the Public Health Agency and is based in Queen's University, Belfast.



INCIDENCE


- There were 814 cases of liver cancer diagnosed during 2018-2022 in Northern Ireland. On average this was 163 cases per year.
- During this period 30.5% of liver cancer cases were among women (Male cases: 566, Female cases: 248). On average there were 113 male and 50 female cases of liver cancer per year.
- The most common diagnosis month during 2018-2022 was January and July among males with 12 cases per year and January, December, May and September among females with 5 cases per year.

Figure 1: Average number of cases of liver cancer per year in 2018-2022 by month of diagnosis



- Liver cancer made up 2.1% of all male and 1.0% of all female cancer cases (excluding non-melanoma skin cancer).
- The liver cancer incidence rates for each gender were 12.1 cases per 100,000 males and 5.1 cases per 100,000 females.
- The odds of developing liver cancer before age 85 was 1 in 73 for men and 1 in 195 for women.

Figure 2: Odds of developing liver cancer in 2018-2022



INCIDENCE BY AGE

- The median age of patients diagnosed with liver cancer during 2018-2022 was 72 years (Males: 72, Females: 73).
- The risk of developing liver cancer varied by age, with 38.2% of men and 47.2% of women diagnosed with liver cancer aged 75 and over at diagnosis.
- In contrast, 8.2% of patients diagnosed with liver cancer were aged 0 to 54 at diagnosis.

Figure 3: Average number of cases of liver cancer diagnosed per year in 2018-2022 by age at diagnosis






Figure 4: Age-specific incidence rates of liver cancer in 2018-2022



INCIDENCE TRENDS


- The number of cases of liver cancer among males increased between 2013-2017 and 2018-2022 by 31.3% from 431 cases (86 cases per year) to 566 cases (113 cases per year).
- The number of cases of liver cancer among females increased between 2013-2017 and 2018-2022 by 3.8% from 239 cases (48 cases per year) to 248 cases (50 cases per year).

Figure 5: Trends in number of cases of liver cancer diagnosed from 2003 to 2022



- Male age-standardised liver cancer incidence rates increased between 2013-2017 and 2018-2022 by 15.2% from 12.5 to 14.4 cases per 100,000 males. This change was not statistically significant.
- Female age-standardised liver cancer incidence rates decreased between 2013-2017 and 2018-2022 by 3.6% from 5.6 to 5.4 cases per 100,000 females. This change was not statistically significant.

Figure 6: Trends in incidence rates of liver cancer from 2003 to 2022



Age-standardised incidence rates illustrate the change in the number of cases within a population of a fixed size and age structure (2013 European Standard).

They thus represent changes other than those caused by population growth and/or ageing.

Trends can also be influenced by changes in how cancer is classified and coded. (e.g. the move from ICD-0-2 to ICD-0-3 in 2019).


INCIDENCE BY STAGE AT DIAGNOSIS

- During 2018-2022 82.1% of liver cancer cases had a stage assigned.
- 21.1% of liver cancer cases were diagnosed at Stage I. (25.7% of staged cases)
- 24.0% of liver cancer cases were diagnosed at Stage IV. (29.2% of staged cases)

Table 1: Number of cases of liver cancer diagnosed in 2018-2022 by stage at diagnosis

Stage at diagnosis	All persons		Male		Female	
	Total cases in period	Average cases per year	Total cases in period	Average cases per year	Total cases in period	Average cases per year
All stages	814	163	566	113	248	50
Stage I	172	34	129	26	43	9
Stage II	125	25	92	18	33	7
Stage III	176	35	132	26	44	9
Stage IV	195	39	120	24	75	15
Unknown	146	29	93	19	53	11

Figure 7: Proportion of cases of liver cancer diagnosed in 2018-2022 by stage at diagnosis




INCIDENCE BY DEPRIVATION

- The number of cases of liver cancer diagnosed during 2018-2022 varied in each deprivation quintile due to variations in population size and age.
- After accounting for these factors, incidence rates:
 - in the most socio-economically deprived areas were 32.9% higher than the NI average.
 - in the least socio-economically deprived areas did not vary significantly from the NI average.

Table 2: Number of cases of liver cancer diagnosed in 2018-2022 by deprivation quintile

Deprivation quintile	All persons		Male		Female	
	Total cases in period	Average cases per year	Total cases in period	Average cases per year	Total cases in period	Average cases per year
Northern Ireland	814	163	566	113	248	50
Most deprived
Quintile 2	174	35	126	25	48	10
Quintile 3	169	34	124	25	45	9
Quintile 4	154	31	97	19	57	11
Least deprived	161	32	110	22	51	10
Unknown	156	31	109	22	47	9
Unknown	0	0	0	0	0	0

Figure 8: Standardised incidence ratio comparing deprivation quintile to Northern Ireland for liver cancer diagnosed in 2018-2022




INCIDENCE BY HEALTH AND SOCIAL CARE TRUST

- The number of cases of liver cancer diagnosed during 2018-2022 varied in each Health and Social Care Trust due to variations in population size and age.
- After accounting for these factors, incidence rates:
 - in Belfast HSCT were 26.6% higher than the NI average.
 - in Northern HSCT did not vary significantly from the NI average.
 - in South Eastern HSCT did not vary significantly from the NI average.
 - in Southern HSCT did not vary significantly from the NI average.
 - in Western HSCT did not vary significantly from the NI average.

Table 3: Number of cases of liver cancer diagnosed in 2018-2022 by Health and Social Care Trust

Health and Social Care Trust	All persons		Male		Female	
	Total cases in period	Average cases per year	Total cases in period	Average cases per year	Total cases in period	Average cases per year
Northern Ireland	814	163	566	113	248	50
Belfast HSCT	182	36	124	25	58	12
Northern HSCT	191	38	136	27	55	11
South Eastern HSCT	191	38	132	26	59	12
Southern HSCT	134	27	89	18	45	9
Western HSCT	116	23	85	17	31	6
Unknown	0	0	0	0	0	0

Figure 9: Standardised incidence ratio comparing Health and Social Care Trust to Northern Ireland for liver cancer diagnosed in 2018-2022




SURVIVAL

- 35.7% of patients were alive one year and 10.3% were alive five years from a liver cancer diagnosis in 2013-2017. (observed survival)
- Age-standardised net survival (ASNS), which removes the effect of deaths from causes unrelated to cancer, was 40.2% one year and 13.5% five years from a liver cancer diagnosis in 2013-2017.
- Five-year survival (ASNS) for liver cancer patients diagnosed in 2013-2017 was 15.3% among men and 10.2% among women.

Table 4: Survival from liver cancer for patients diagnosed in 2013-2017

Time since diagnosis	All persons		Male		Female	
	Observed survival	Age-standardised net survival	Observed survival	Age-standardised net survival	Observed survival	Age-standardised net survival
6 months	48.7%	52.0%	50.8%	52.4%	44.9%	51.5%
One year	35.7%	40.2%	39.3%	41.3%	29.2%	38.3%
Two years	22.5%	26.2%	25.5%	27.5%	16.9%	23.8%
Five years	10.3%	13.5%	12.2%	15.3%	6.8%	10.2%

Figure 10: Age-standardised net survival from liver cancer for patients diagnosed in 2013-2017



Observed survival examines the time between diagnosis and death from any cause, however, due to the inclusion of non-cancer deaths it may not fully reflect how changes in cancer care impact survival from cancer.

Age-standardised net survival provides an estimate of patient survival which has been adjusted to take account of deaths unrelated to cancer. It is more widely used to assess the impact of changes in cancer care on patient survival.


SURVIVAL BY AGE

- Survival from liver cancer among patients diagnosed during 2013-2017 was related to age with better five-year survival among younger age groups.
- Five-year net survival ranged from 20.3% among patients aged 55 to 64 at diagnosis to 3.9% among those aged 75 to 99.

Table 5: Net survival from liver cancer for patients diagnosed in 2013-2017 by age at diagnosis

Age group	All persons	
	One-year	Five-years
15 to 54	41.5%	17.2%
55 to 64	50.4%	20.3%
65 to 74	45.8%	16.7%
75 to 99	25.2%	3.9%

Figure 11: Net survival from liver cancer for patients diagnosed in 2013-2017 by age at diagnosis




SURVIVAL TRENDS

ONE-YEAR NET SURVIVAL

- Between 2013-2017 and 2018-2022 there was no significant change in one-year survival (ASNS) from liver cancer.
- Compared to 1993-1997 one-year survival (ASNS) from liver cancer in 2018-2022 increased significantly from 18.6% to 40.7%. This increase was significant for males (22.6% to 42.7%) and females (14.2% to 36.0%).


Figure 12: Trends in one-year age-standardised net survival from liver cancer in 1993-2022



FIVE-YEAR NET SURVIVAL

- Between 2008-2012 and 2013-2017 there was no significant change in five-year survival (ASNS) from liver cancer.
- Compared to 1993-1997 five-year survival (ASNS) from liver cancer in 2013-2017 increased significantly from 4.1% to 13.5%. This increase was significant for males (3.9% to 15.3%) but not females.

Figure 13: Trends in five-year age-standardised net survival from liver cancer in 1993-2017




SURVIVAL BY STAGE

- Survival from liver cancer among patients diagnosed during 2013-2017 was strongly related to stage with better five-year survival among those diagnosed at earlier stages.
- Five-year survival (ASNS) ranged from 48.3% among patients diagnosed at Stage I to 1.6% among those diagnosed at Stage IV.

Table 6: Age-standardised net survival from liver cancer for patients diagnosed in 2013-2017 by stage at diagnosis

Stage at diagnosis	All persons	
	One-year	Five-years
Stage I	80.2%	48.3%
Stage II	61.6%	14.5%
Stage III	27.6%	3.2%
Stage IV	19.0%	1.6%
Unknown	34.4%	10.4%

Figure 14: Age-standardised net survival from liver cancer for patients diagnosed in 2013-2017 by stage at diagnosis



PREVALENCE

- At the end of 2022, there were 312 people (Males: 226; Females: 86) living with liver cancer who had been diagnosed with the disease during 1998-2022.
- Of these 31.7% had been diagnosed in the previous year (one-year prevalence) and 85.9% in the previous 10 years (ten-year prevalence).
- 35.3% of liver cancer survivors were aged 75 and over at the end of 2022.


Table 7: 25-year prevalence of liver cancer by age at end of 2022

Gender	Age at end of 2022	25-year prevalence	Time since diagnosis			
			0 to 1 year	1 to 5 years	5 to 10 years	10 to 25 years
All persons	All ages	312	99	115	54	44
	0 to 74	202	63	72	35	32
	75 and over	110	36	43	19	12
Male	All ages	226	76	82	41	27
	0 to 74	144	47	49	28	20
	75 and over	82	29	33	13	7
Female	All ages	86	23	33	13	17
	0 to 74	58	16	23	7	12
	75 and over	28	7	10	6	5

PREVALENCE TRENDS

- 10-year prevalence of liver cancer among males increased between 2017 and 2022 by 33.6% from 149 survivors to 199 survivors.
- 10-year prevalence of liver cancer among females decreased between 2017 and 2022 by 1.4% from 70 survivors to 69 survivors.

Figure 15: Trends in 10-year prevalence of liver cancer in 2002-2022



MORTALITY

- There were 875 deaths from liver cancer during 2018-2022 in Northern Ireland. On average this was 175 deaths per year.
- During this period 40.6% of liver cancer deaths were among women (Male deaths: 520, Female deaths: 355). On average there were 104 male and 71 female deaths from liver cancer per year.
- Liver cancer deaths made up 4.3% of all male and 3.3% of all female cancer deaths.
- The median age of patients who died from liver cancer during 2018-2022 was 75 years (Males: 74, Females: 77).
- The risk of dying from liver cancer varied by age, with 48.7% of men and 59.7% of women who died from liver cancer aged 75 and over at death.
- In contrast, 5.3% of patients who died from liver cancer were aged 0 to 54 at death.

Figure 16: Average number of deaths from liver cancer per year in 2018-2022 by age at death






Figure 17: Age-specific mortality rates of liver cancer in 2018-2022



MORTALITY TRENDS


- The number of deaths from liver cancer among males increased between 2013-2017 and 2018-2022 by 21.8% from 427 deaths (85 deaths per year) to 520 deaths (104 deaths per year).
- The number of deaths from liver cancer among females increased between 2013-2017 and 2018-2022 by 23.3% from 288 deaths (58 deaths per year) to 355 deaths (71 deaths per year).

Figure 18: Trends in the number of deaths from liver cancer from 2003 to 2022



- Male age-standardised liver cancer mortality rates increased between 2013-2017 and 2018-2022 by 6.2% from 12.9 to 13.7 deaths per 100,000 males. This change was not statistically significant.
- Female age-standardised liver cancer mortality rates increased between 2013-2017 and 2018-2022 by 11.8% from 6.8 to 7.6 deaths per 100,000 females. This change was not statistically significant.

Figure 19: Trends in mortality rates of liver cancer from 2003 to 2022



Age-standardised mortality rates illustrate the change in the number of deaths within a population of a fixed size and age structure (2013 European Standard).

They thus represent changes other than those caused by population growth and/or ageing.

Trends can also be influenced by changes in how cancer is classified and coded.

BACKGROUND NOTES

Cancer classification: Classification of tumour sites is carried out using ICD10 codes. For a listing and explanation of ICD10 codes see: World Health Organisation at <http://apps.who.int/classifications/icd10/browse/2010/en#/I>

Population data: Population data for Northern Ireland, and smaller geographic areas, are extracted from the NI mid-year population estimates available from the NI Statistics and Research Agency (available at www.nisra.gov.uk).

Geographic areas: Geographic areas are assigned based on a patient's postcode of usual residence at diagnosis using the Jul 2024 Central Postcode Directory (CPD) produced by the NI Statistics and Research Agency (available at www.nisra.gov.uk).

Deprivation quintiles: Super output areas (SOA) are assigned to each patient based on their postcode of usual residence at diagnosis. Using the SOA each patient is assigned a socio-economic deprivation quintile based on the 2017 Multiple Deprivation Measure. The 2017 Multiple Deprivation Measure is available from the NI Statistics and Research Agency (available at www.nisra.gov.uk).

Crude incidence/mortality rate: The number of cases/deaths per 100,000 person years in the population. Person years are the sum of the population over the number of years included.

Age-standardised incidence/mortality rates per 100,000 person years are estimates of the incidence/mortality rate if that population had a standard age structure. Throughout this report the 2013 European Standard Population has been used. Standardising to a common Standard Population allows comparisons of incidence/mortality rates to be made between different time periods and geographic areas while removing the effects of population change and ageing.

Standardised Incidence/Mortality Ratio (SIR/SMR) is the ratio of the number of cases/deaths observed in a population to the expected number of cases/deaths, based upon the age-specific rates in a reference population. This statistic is often used to compare incidence/mortality rates for geographic areas (e.g. Trusts) to the national incidence/mortality rates (i.e. Northern Ireland). An SIR/SMR of 100 indicates there is no difference between the geographic area and the national average.

Confidence intervals measure the precision of a statistic (e.g. liver cancer incidence rate). Typically, when numbers are low, precision is poorer and confidence intervals will be wider. As a general rule, when comparing statistics (e.g. liver cancer incidence rate in year 2012 vs year 2013), if the confidence interval around one statistic overlaps with the interval around another, it is unlikely that there is any real difference between the two. If there is no overlap, the difference is considered to be statistically significant.

Lifetime risk is estimated as the cumulative risk of getting cancer up to age 75/85, calculated directly from the age-specific incidence rates. The odds of developing the disease before age 75/85 is the inverse of the cumulative risk.

Prevalence is the number of cancer patients who are alive in the population on a specific date (31st December 2022 in this report). Since data from the NI Cancer Registry are only available since 1993, prevalence only refers to a fixed term (10 and 25 years in this report). There may be members of the population living with a diagnosis of cancer for more than 25 years.

Patient survival is evaluated using two measures. Observed survival examines the time between diagnosis and death from any cause. It thus represents what cancer patients experience, however, due to the inclusion of non-cancer deaths (e.g. heart disease), it may not reflect how changes in cancer care impact survival from cancer. Thus age-standardised net survival is also examined. This measure provides an estimate of patient survival which has been adjusted to take account of deaths unrelated to cancer. It also assumes a standard age distribution thereby removing the impact of changes in the age distribution of cancer patients on changes in survival over time. While this measure is hypothetical, as it assumes patients can only die from cancer related factors, it is a better indicator of the impact of changes in cancer care on patient survival.