

Cancer among older people

(excluding non-melanoma skin cancer)

Patients diagnosed 1993-2019
(Ages 75 and over; ICD10: C00-C43, C45-C97)

Further information

Further data is available at: www.qub.ac.uk/research-centres/nicr

Phone: +44 (0)28 9097 6028

e-mail: nicr@qub.ac.uk

Acknowledgements

The Northern Ireland Cancer Registry (NICR) is funded by the Public Health Agency and is based in Queen's University, Belfast. NICR uses data provided by patients and collected by the health service as part of their care and support.

The incidence, prevalence and survival statistics in this publication are designated as official statistics signifying that they comply with the Code of Practice for Official Statistics.




Incidence

- During 2015-2019 there were 1,745 male and 1,605 female cases of cancer (ex NMSC) diagnosed among those aged 75 and over each year.
- Cancer (ex NMSC) among those aged 75 and over made up 35.2% of all male, and 32.7% of female cancers (ex NMSC) among all age groups.

Incidence by year of diagnosis - Cancer among older people, Cases in 2010-2019


- The number of cases of cancer (ex NMSC) increased by 14.2% between 2010-2014 and 2015-2019 among older males aged 75 and over, while among older females the number of cases increased by 10.1%.

Year of diagnosis	Male	Female	Both sexes
2010	1,417	1,363	2,780
2011	1,552	1,457	3,009
2012	1,502	1,452	2,954
2013	1,564	1,504	3,068
2014	1,606	1,513	3,119
2015	1,662	1,553	3,215
2016	1,730	1,634	3,364
2017	1,732	1,635	3,367
2018	1,801	1,592	3,393
2019	1,800	1,612	3,412



Trends in age-standardised incidence rates - Cancer among older people, Cases in 1995-2019

- Among older males age-standardised incidence rates of cancer (ex NMSC) decreased by 2.9% from 3,222.8 per 100,000 person years in 2010-2014 to 3,128.9 cases per 100,000 persons years in 2015-2019. This difference was not statistically significant.
- Among older females age-standardised incidence rates of cancer (ex NMSC) increased by 2.3% from 1,956.9 per 100,000 person years in 2010-2014 to 2,001.2 cases per 100,000 persons years in 2015-2019. This difference was not statistically significant.



Age-standardised incidence rates illustrate the change in the number of cases within a population of a fixed size and age structure (2013 European Standard).

They thus represent changes other than those caused by population growth and/or ageing.


Trends can also be influenced by changes in how cancer is classified and coded. (e.g. the move from ICD-0-2 to ICD-0-3 in 2019).

Note: Annual averages have been rounded to the nearest integer. Sums of numbers in table rows or columns may thus differ slightly from the given total.

NMSC: Non-melanoma skin cancer

Incidence by cancer type - Cancer among older people, Cases in 2015-2019

The most common cancer types among older men (excluding NMSC), were prostate cancer (24.4%), lung cancer (16.5%) and colorectal cancer (14.8%), while the most common cancer types among older women (excluding NMSC) were breast cancer (20.8%), lung cancer (16.5%) and colorectal cancer (14.2%).



Incidence by deprivation quintile - Cancer among older people, Cases in 2015-2019

The annual number of cases during 2015-2019 varied in each deprivation quintile due to variations in population size and age.

After accounting for these factors, incidence rates:


- in the most socio-economically deprived areas were 10.2% higher than the NI average.
- in the least socio-economically deprived areas did not vary significantly from the NI average.

Deprivation quintile	Average cases per year		
	Male	Female	Both sexes
Least deprived (Quintile 1)	389	345	734
Quintile 2	355	319	673
Quintile 3	357	319	676
Quintile 4	356	320	677
Most deprived (Quintile 5)	288	303	590
Northern Ireland	1,745	1,605	3,350

Standardised incidence ratios compare incidence rates in each deprivation quintile with the Northern Ireland incidence rate.

A value above 100 means that incidence rates in that deprivation quintile are greater than the Northern Ireland average.

This measure takes account of population size and age structure. Differences are thus not a result of these factors.




Incidence by Health and Social Care Trust (HSCT) - Cancer among older people, Cases in 2015-2019

The annual number of cases during 2015-2019 varied in each HSCT due to variations in population size and age.

After accounting for these factors, incidence rates:

- in Belfast HSCT were significantly higher than the NI average.
- in Northern HSCT did not vary significantly from the NI average.
- in South-Eastern HSCT did not vary significantly from the NI average.
- in Southern HSCT did not vary significantly from the NI average.
- in Western HSCT did not vary significantly from the NI average.

Health and Social Care Trust	Average cases per year		
	Male	Female	Both sexes
Belfast HSCT	345	362	706
Northern HSCT	462	423	885
South-Eastern HSCT	367	311	678
Southern HSCT	312	281	593
Western HSCT	258	229	487
Northern Ireland	1,745	1,605	3,350



Standardised incidence ratios compare incidence rates in each HSC Trust with the Northern Ireland incidence rate.

A value above 100 means that incidence rates in that HSC Trust are greater than the NI average.

This measure takes account of population size and age structure. Differences are thus not a result of these factors.


Data for Local Government Districts and Parliamentary Constituencies are available at www.qub.ac.uk/research-centres/nicr

Incidence by method of most recent admission to hospital - Cancer among older people, Cases in 2015-2019

During 2015-2019:

- 22.7% of cases had an emergency admission to hospital recorded within 30 days prior to their cancer diagnosis.
- 21.9% of male cases had an emergency admission up to 30 days prior to diagnosis, compared to 23.5% of female cases.
- In 44.6% of diagnosed cases there was no record of a hospital inpatient admission up to 30 days prior to the diagnosis.

Method of admission	Average cases per year		
	Male	Female	Both sexes
Emergency admission	383	378	760
Elective admission	661	433	1,095
No emergency/elective admission recorded	701	794	1,495
Total	1,745	1,605	3,350



Admission method refers to the most recent hospital inpatient admission that a patient had prior to cancer diagnosis, regardless of reason for the admission.

Admissions are considered up to a maximum of 30 days prior to diagnosis. Admissions up to two days post diagnosis are also considered to allow for a reasonable margin or error in data recording.


The majority of patients with no inpatient admission recorded prior to diagnosis are likely to have been diagnosed via an outpatient route.

Incidence by stage at diagnosis - Cancer among older people, Cases in 2015-2019

During 2015-2019:

- 74.2% of cases diagnosed had a stage assigned.
- 20.3% of cases were diagnosed at stage I. (27.3% of staged cases)
- 23.0% of cases were diagnosed at stage IV. (31.0% of staged cases)
- Among cases which were staged, 34.5% of male cases were diagnosed at stage IV, compared to 27.2% of female cases.

Stage at diagnosis	Average cases per year		
	Male	Female	Both sexes
Stage I (Early)	338	341	679
Stage II	216	295	510
Stage III	294	231	525
Stage IV (Late)	447	324	771
Unknown	450	415	865
All stages	1,745	1,605	3,350



Cancer stage describes the size of a cancer and how far it has grown and spread. This information is used to help decide what treatments are needed. The classification used here to stage cancer is the TNM classification (Version 7 prior to 2018, Version 8 from 2018 onwards).


Survival

- 53.5% of older patients were alive one year and 26.5% were alive five years from a cancer (ex NMSC) diagnosis in 2010-2014. (observed survival)
- Net survival, which removes the effect of deaths from causes unrelated to cancer, was 57.7% one year and 40.1% five years from a cancer (ex NMSC) diagnosis among those aged 75 and over in 2010-2014.
- Five-year net survival for older patients diagnosed in 2010-2014 was 41.2% among men and 39.1% among women.

Gender	Observed survival		Net survival	
	One-year	Five-years	One-year	Five-years
Male	55.0%	26.4%	59.7%	41.2%
Female	52.0%	26.7%	55.7%	39.1%
Both sexes	53.5%	26.5%	57.7%	40.1%


Observed survival is the proportion of patients still alive one/five years after diagnosis. However, in this measure patients may have died from causes unrelated to their cancer.

Net survival is the proportion of patients who would survive if the patient could not die from causes unrelated to their cancer. This measure is more typically used in studies of cancer survival.




Trends in survival - Cancer among older people, Patients diagnosed in 1995-2014

- Among men five-year survival from cancer (ex NMSC) among those aged 75 and over decreased from 43.0% in 2005-2009 to 41.2% in 2010-2014. This difference was not statistically significant.
- Among women five-year survival from cancer (ex NMSC) among those aged 75 and over increased from 35.9% in 2005-2009 to 39.1% in 2010-2014. This difference was not statistically significant.




Survival by cancer type - Cancer among older people, Patients diagnosed in 2010-2014

- Five-year survival for older patients diagnosed in 2010-2014 ranged from 80.5% for melanoma to 1.3% for pancreatic cancer.
- In particular five-year survival for the most common cancer types among older people was 73.7% for female breast cancer, 50.9% for colorectal cancer, 6.4% for lung cancer and 74.3% for prostate cancer.



Survival by stage at diagnosis - Cancer among older people, Patients diagnosed in 2010-2014


- Stage at diagnosis is one of the most important factors in cancer (ex NMSC) survival among older people.
- Five-year survival ranged from 82.8% for early stage (stage I) disease to 8.2% for late stage (stage IV) disease.
- Five-year survival (ASNS) for unstaged cancer was 22.9%.
- Five-year survival (ASNS) among older people for stage IV cancer was 10.4% for men, compared to 5.4% for women.



Survival by method of most recent admission to hospital - Cancer among older people, Patients diagnosed in 2010-2014

Five-year survival among older patients who had an emergency admission to hospital up to 30 days prior to their cancer diagnosis was 11.1% compared to 54.7% among those with elective admissions and 48.0% among those who had neither type of hospital admission recorded up to 30 days prior to diagnosis.

Five-year survival (ASNS) among older patients who had an emergency admission to hospital within 30 days prior to their cancer diagnosis was 10.5% for men, compared to 11.6% for women.



Prevalence

At the end of 2019, there were 23,026 people aged 75 and over (Males: 11,500; Females: 11,526) living with cancer (ex NMSC) who had been diagnosed with the disease during 1995-2019.


Of these, 49.9% were male, and 10.4% had been diagnosed in the previous year.

Time since diagnosis	Male	Female	Both sexes
0-1 year	1,307	1,092	2,399
1-5 years	3,537	2,910	6,447
5-10 years	2,943	2,855	5,798
10-25 years	3,713	4,669	8,382
0-25 years	11,500	11,526	23,026

25-year prevalence refers to the number of cancer survivors who were alive at the end of 2019, and had been diagnosed with their cancer in the previous 25 years (i.e. 1995-2019).

25-year prevalence by cancer type - Cancer among older people, Patients alive at end of 2019

The most prevalent cancer types among male survivors aged 75 and over at the end of 2019 (ex NMSC), were prostate cancer (5,657 survivors) and colorectal cancer (2,273 survivors), while the most prevalent cancer types among female survivors aged 75 and over were breast cancer (4,787 survivors) and colorectal cancer (2,085 survivors).



Trends in 10-year prevalence - Cancer among older people, Patients alive at end of each year from 2010-2019

- Among older males the number of survivors from cancer (ex NMSC) who had been diagnosed within the previous five years increased by 18.6% from 6,566 survivors in 2014 to 7,787 survivors in 2019.
- Among older females the number of survivors from cancer (ex NMSC) who had been diagnosed within the previous five years increased by 22.3% from 5,608 survivors in 2014 to 6,857 survivors in 2019.

	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
Male	5,341	5,685	5,961	6,296	6,566	6,730	6,943	7,153	7,565	7,787
Female	4,761	5,050	5,243	5,432	5,608	5,915	6,181	6,436	6,666	6,857
Both sexes	10,102	10,735	11,204	11,728	12,174	12,645	13,124	13,589	14,231	14,644

Mortality

During 2015-2019 there were 1,173 male and 1,085 female deaths among older people (aged 75 and over) from cancer (ex NMSC) each year.


Deaths by year of death - Cancer among older people, Deaths in 2010-2019

- Among males the number of deaths from cancer (ex NMSC) increased by 15.3% from an annual average of 1,017 deaths in 2010-2014 to 1,173 deaths in 2015-2019.
- Among females the number of deaths from cancer (ex NMSC) increased by 6.8% from an annual average of 1,016 deaths in 2010-2014 to 1,085 deaths in 2015-2019.

	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
Male	950	1,048	1,007	1,032	1,046	1,169	1,142	1,152	1,172	1,232
Female	951	969	1,011	1,058	1,093	988	1,109	1,097	1,106	1,127
Both sexes	1,901	2,017	2,018	2,090	2,139	2,157	2,251	2,249	2,278	2,359

Trends in age-standardised mortality rates - All cancers (ex NMSC), Deaths in 1995-2019

- Among males aged 75 and over age-standardised mortality rates from cancer (ex NMSC) decreased by 0.8% between 2010-2014 and 2015-2019 from 2,222.2 to 2,204.8 deaths per 100,000 persons years. This difference was not statistically significant.
- Among females aged 75 and over age-standardised mortality rates from cancer (ex NMSC) decreased by 1.5% between 2010-2014 and 2015-2019 from 1,352.6 to 1,332.1 deaths per 100,000 persons years. This difference was not statistically significant.




Mortality data are provided by the Northern Ireland General Registrar Office via the Department of Health.

Counts of the number of deaths are based upon the year that death occurred, and upon the primary cause of death only.

Age-standardised mortality rates remove changes over time caused by population growth and/or ageing.

Deaths by cancer type - Cancer among older people, Deaths in 2015-2019

The most common causes of cancer death (ex NMSC) among men, were lung cancer (22.2%), prostate cancer (17.5%) and colorectal cancer (11.0%), while the most common cause of cancer death (ex NMSC) among women were lung cancer (19.9%), breast cancer (13.5%) and colorectal cancer (11.7%).




Deaths by deprivation quintile - Cancer among older people, Deaths in 2015-2019

The annual number of deaths during 2015-2019 varied in each deprivation quintile due to variations in population size and age.

After accounting for these factors, mortality rates:

- in the most socio-economically deprived areas were 20.4% higher than the NI average.
- in the least socio-economically deprived areas were 9.2% lower than the NI average.

Deprivation quintile	Average deaths per year		
	Male	Female	Both sexes
Least deprived (Quintile 1)	240	218	458
Quintile 2	238	203	441
Quintile 3	244	221	465
Quintile 4	240	224	463
Most deprived (Quintile 5)	211	219	430
Northern Ireland	1,173	1,085	2,259



Standardised mortality ratios compare mortality rates in each deprivation quintile with the Northern Ireland mortality rate.

A value above 100 means that mortality rates in that deprivation quintile are greater than the Northern Ireland average.

This measure takes account of population size and age structure. Differences are thus not a result of these factors.

Background notes

Cancer classification: Classification of tumour sites is carried out using ICD10 codes. For a listing and explanation of ICD10 codes see: World Health Organisation at <http://apps.who.int/classifications/icd10/browse/2010/en#/II>

Population data for Northern Ireland, and smaller geographic areas, are extracted from the NI mid-year population estimates available from the NI Statistics and Research Agency (available at www.nisra.gov.uk).

Geographic areas are assigned based on a patient's postcode of usual residence at diagnosis using the Jan 2021 Central Postcode Directory (CPD) produced by the NI Statistics and Research Agency (available at www.nisra.gov.uk).

Deprivation quintiles: Super output areas (SOA) are assigned to each patient based on their postcode of usual residence at diagnosis. Using the SOA each patient is assigned a socio-economic deprivation quintile based on the 2017 Multiple Deprivation Measure. The 2017 Multiple Deprivation Measure is available from the NI Statistics and Research Agency (available at www.nisra.gov.uk).

A **crude incidence/mortality rate** is the number of cases/deaths per 100,000 person years in the population. Person years are the sum of the population over the number of years included.

An **age-standardised incidence/mortality rate** per 100,000 person years is an estimate of the incidence/mortality rate if that population had a standard age structure. Throughout this report the 2013 European Standard Population has been used. Standardising to a common Standard Population allows comparisons of incidence/mortality rates to be made between different time periods and geographic areas while removing the effects of population change and ageing.

A **Standardised Incidence/Mortality Ratio (SIR/SMR)** is the ratio of the number of cases/deaths observed in a population to the expected number of cases/deaths, based upon the age-specific rates in a reference population. This statistic is often used to compare incidence/mortality rates for geographic areas (e.g. Trusts) to the national incidence/mortality rates (i.e. Northern Ireland). An SIR/SMR of 100 indicates there is no difference between the geographic area and the national

Confidence intervals are a measure of the precision of a statistic (e.g. colorectal cancer incidence rate). Typically, when numbers are low, precision is poorer and confidence intervals will be wider. As a general rule, when comparing statistics (e.g. cervical cancer incidence rate in year 2012 vs year 2013), if the confidence interval around one statistic overlaps with the interval around another, it is unlikely that there is any real difference between the two. If there is no overlap, the difference is considered to be **statistically significant**.

Lifetime risk is estimated as the cumulative risk of getting cancer up to age 75/85, calculated directly from the age-specific incidence rates. The odds of developing the disease before age 75/85 is the inverse of the cumulative risk.

Prevalence is the number of cancer patients who are alive in the population on a specific date (31st December 2019 in this report). Since data from the NI Cancer Registry are only available since 1993, prevalence only refers to a fixed term (10 and 25 years in this report). There may be members of the population living with a diagnosis of cancer for more than 25 years.

Observed survival refers to the proportion of patients who survive a specified amount of time from their date of diagnosis. Observed survival considers death from any cause and is not adjusted for the age of the patient. Cause of death may be unrelated to the cancer the patient has been diagnosed with.

Net Survival is an estimate of survival where the effect on survival of background population mortality rates has been removed. It represents the [theoretical] survival of cancer patients if they could only die from cancer-related causes. Age-standardised net survival estimates are the estimates that would occur if that population of cancer patients had a standard population age structure. The age groups and weights used here are those used by international studies such as EUROCORE, an international study group that compares cancer survival among European countries. However, due to the small number of patients in NI, the first two age categories in the standard population are combined.

Mortality: Information relating to cancer mortality is sourced from the General Registrar Office (GRONI) via the Department of Health (NI). Results are based upon the date on which death occurs, and may thus differ slightly than those produced by the Northern Ireland Statistics and Research Agency (NISRA), which produces deaths data based upon the date on which the death is registered with GRONI.