
Breast insitu tumours

Patients diagnosed 1993-2020
(ICD10: D05)

Further information

Further data is available at: www.qub.ac.uk/research-centres/nicr

Phone: +44 (0)28 9097 6028

e-mail: nicr@qub.ac.uk

Acknowledgements

The Northern Ireland Cancer Registry (NICR) is funded by the Public Health Agency and is based in Queen's University, Belfast. NICR uses data provided by patients and collected by the health service as part of their care and support.

The incidence, prevalence and survival statistics in this publication are designated as official statistics signifying that they comply with the Code of Practice for Official Statistics.



Incidence

During 2016-2020:


- There were 240 female breast insitu tumours diagnosed each year.
- The risk of being diagnosed with a breast insitu tumour before the age of 75 was 1 in 52 for women, while before the age of 85 the risk was 1 in 45.

Incidence by age at diagnosis - Breast insitu tumours, Cases in 2016-2020

During 2016-2020:

- The median age at diagnosis among women was 57.
- Cancer risk varied with age, with 9.6% of women aged 75 years or more at diagnosis, compared to 42.9% diagnosed among those aged under 55.


Age at diagnosis	Average cases per year
	Female
0 - 54	103
55 - 64	61
65 - 74	54
75 +	23
All ages	240



Incidence by year of diagnosis - Breast insitu tumours, Cases in 1996-2020

- The number of female breast insitu tumours increased by 41.2% from an annual average of 170 cases in 2011-2015 to 240 cases in 2016-2020.


Year of diagnosis	Female cases
2011	163
2012	129
2013	155
2014	161
2015	240
2016	250
2017	241
2018	297
2019	216
2020	194



Note: Annual averages based upon several years have been rounded to the nearest integer. Sums of numbers in table rows or columns may thus differ slightly from the given total.

Trends in age-standardised incidence rates - Breast insitu tumours, Cases in 1996-2020

Among females the age-standardised incidence rate of breast insitu tumours increased by 32.8% from 20.1 per 100,000 person years in 2011-2015 to 26.7 cases per 100,000 persons years in 2016-2020. This difference was statistically significant.



Age-standardised incidence rates illustrate the change in the number of cases within a population of a fixed size and age structure (2013 European Standard).

They thus represent changes other than those caused by population growth and/or ageing.

Trends can also be influenced by changes in how cancer is classified and coded. (e.g. the move from ICD-0-2 to ICD-0-3 in 2019).

Incidence by deprivation quintile - Breast insitu tumours, Cases in 2016-2020

The annual number of cases during 2016-2020 varied in each deprivation quintile due to variations in population size and age.

After accounting for these factors, incidence rates:


- in the least socio-economically deprived areas were 16.5% higher than the NI average.
- in the most socio-economically deprived areas were 15.7% lower than the NI average.

Deprivation quintile	Average cases per year
	Female
Most deprived (Quintile 1)	35
Quintile 2	47
Quintile 3	48
Quintile 4	52
Least deprived (Quintile 5)	58
Northern Ireland	240

Standardised incidence ratios compare incidence rates in each deprivation quintile with the Northern Ireland incidence rate.

A value above 100 means that incidence rates in that deprivation quintile are greater than the Northern Ireland average.

This measure takes account of population size and age structure. Differences are thus not a result of these factors.




Incidence by Health and Social Care Trust (HSCT) - Breast insitu tumours, Cases in 2016-2020

The annual number of cases during 2016-2020 varied in each HSCT due to variations in population size and age.

After accounting for these factors, incidence rates:

- in Belfast HSCT did not vary significantly from the NI average.
- in Northern HSCT did not vary significantly from the NI average.
- in South-Eastern HSCT did not vary significantly from the NI average.
- in Southern HSCT did not vary significantly from the NI average.
- in Western HSCT did not vary significantly from the NI average.

Health and Social Care Trust	Average cases per year
	Female
Belfast HSCT	42
Northern HSCT	69
South Eastern HSCT	46
Southern HSCT	41
Western HSCT	41
Northern Ireland	240



Standardised incidence ratios compare incidence rates in each HSC Trust with the Northern Ireland incidence rate. A value above 100 means that incidence rates in that HSC Trust are greater than the NI average.

This measure takes account of population size and age structure. Differences are thus not a result of these factors.

Data for Local Government Districts and Parliamentary Constituencies are available at www.qub.ac.uk/research-centres/nicr

Prevalence

- At the end of 2020, there were 2,934 women living with breast insitu tumours who had been diagnosed in the previous 25 years.
- Of these, 20.9% were aged 75 and over, and 6.4% had been diagnosed in the previous year.

25-year prevalence refers to the number of cancer survivors who were alive at the end of 2020, and had been diagnosed with their cancer in the previous 25 years (i.e. 1996-2020).

Time since diagnosis	25-year prevalence		
	Aged 0-74	Aged 75+	All ages
0-1 year	164	23	187
1-5 years	815	112	927
5-10 years	589	148	737
10-25 years	753	330	1,083
0-25 years	2,321	613	2,934

Trends in 10-year prevalence - Breast insitu tumours, Patients alive at end of each year from 2011-2020

- Among women the number of survivors from breast insitu tumours who had been diagnosed within the previous ten years increased by 32.4% from 1,398 survivors in 2015 to 1,851 survivors in 2020.

	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
Female	1,121	1,169	1,219	1,280	1,398	1,531	1,614	1,763	1,821	1,851

Background notes

Cancer classification: Classification of tumour sites is carried out using ICD10 codes. For a listing and explanation of ICD10 codes see: World Health Organisation at <http://apps.who.int/classifications/icd10/browse/2010/en#/I>

Population data for Northern Ireland, and smaller geographic areas, are extracted from the NI mid-year population estimates available from the NI Statistics and Research Agency (available at www.nisra.gov.uk).

Geographic areas are assigned based on a patient's postcode of usual residence at diagnosis using the Jan 2021 Central Postcode Directory (CPD) produced by the NI Statistics and Research Agency (available at www.nisra.gov.uk).

Deprivation quintiles: Super output areas (SOA) are assigned to each patient based on their postcode of usual residence at diagnosis. Using the SOA each patient is assigned a socio-economic deprivation quintile based on the 2017 Multiple Deprivation Measure. The 2017 Multiple Deprivation Measure is available from the NI Statistics and Research Agency (available at www.nisra.gov.uk).

A **crude incidence/mortality rate** is the number of cases/deaths per 100,000 person years in the population. Person years are the sum of the population over the number of years included.

An **age-standardised incidence/mortality rate** per 100,000 person years is an estimate of the incidence/mortality rate if that population had a standard age structure. Throughout this report the 2013 European Standard Population has been used. Standardising to a common Standard Population allows comparisons of incidence/mortality rates to be made between different time periods and geographic areas while removing the effects of population change and ageing.

A **Standardised Incidence/Mortality Ratio (SIR/SMR)** is the ratio of the number of cases/deaths observed in a population to the expected number of cases/deaths, based upon the age-specific rates in a reference population. This statistic is often used to compare incidence/mortality rates for geographic areas (e.g. Trusts) to the national incidence/mortality rates (i.e. Northern Ireland). An SIR/SMR of 100 indicates there is no difference between the geographic area and the national average.

Confidence intervals are a measure of the precision of a statistic (e.g. breast insitu tumours incidence rate). Typically, when numbers are low, precision is poorer and confidence intervals will be wider. As a general rule, when comparing statistics (e.g. breast insitu tumours incidence rate in year 2012 vs year 2013), if the confidence interval around one statistic overlaps with the interval around another, it is unlikely that there is any real difference between the two. If there is no overlap, the difference is considered to be **statistically significant**.

Lifetime risk is estimated as the cumulative risk of getting cancer up to age 75/85, calculated directly from the age-specific incidence rates. The odds of developing the disease before age 75/85 is the inverse of the cumulative risk.

Prevalence is the number of cancer patients who are alive in the population on a specific date (31st December 2020 in this report). Since data from the NI Cancer Registry are only available since 1993, prevalence only refers to a fixed term (10 and 25 years in this report). There may be members of the population living with a diagnosis of cancer for more than 25 years.

Observed survival refers to the proportion of patients who survive a specified amount of time from their date of diagnosis. Observed survival considers death from any cause and is not adjusted for the age of the patient. Cause of death may be unrelated to the cancer the patient has been diagnosed with.

Net Survival is an estimate of survival where the effect on survival of background population mortality rates has been removed. It represents the [theoretical] survival of cancer patients if they could only die from cancer-related causes. Age-standardised net survival estimates are the estimates that would occur if that population of cancer patients had a standard population age structure. The age groups and weights used here are those used by international studies such as EURO CARE, an international study group that compares cancer survival among European countries. However, due to the small number of patients in NI, the first two age categories in the standard population are combined.

Mortality: Information relating to cancer mortality is sourced from the General Registrar Office (GRONI) via the Department of Health (NI). Results are based upon the date on which death occurs, and may thus differ slightly than those produced by the Northern Ireland Statistics and Research Agency (NISRA), which produces deaths data based upon the date on which the death is registered with GRONI.