

Trachea, Bronchus & Lung (ICD10 C33-C34) - INCIDENCE 1993-2005

| Gender | Year | No of Cases | % of Total | Crude Rate | EASR | EASR Lower | EASR Upper | WASR | WASR Lower | WASR Upper | Cumulative Risk (0-74 yrs) | Odds, 1 in: |
|---------------|------|-------------|------------|------------|------|------------|------------|------|------------|------------|----------------------------|-------------|
| Male | 1993 | 565 | 13.7% | 70.8 | 77.9 | 71.4 | 84.4 | 52.2 | 47.7 | 56.6 | 6.5 | 15.4 |
| | 1994 | 625 | 15.0% | 77.9 | 87.5 | 80.6 | 94.5 | 59.2 | 54.5 | 64.0 | 7.3 | 13.8 |
| | 1995 | 546 | 13.4% | 67.9 | 73.2 | 67.0 | 79.4 | 48.8 | 44.6 | 53.0 | 6.2 | 16.2 |
| | 1996 | 550 | 13.0% | 67.9 | 72.6 | 66.5 | 78.7 | 48.2 | 44.1 | 52.4 | 6.3 | 15.9 |
| | 1997 | 552 | 13.3% | 67.7 | 72.8 | 66.6 | 78.9 | 48.5 | 44.3 | 52.7 | 6.0 | 16.7 |
| | 1998 | 562 | 13.3% | 68.6 | 71.8 | 65.8 | 77.7 | 47.7 | 43.7 | 51.8 | 6.0 | 16.8 |
| | 1999 | 511 | 12.2% | 62.4 | 64.9 | 59.2 | 70.6 | 42.6 | 38.8 | 46.4 | 5.3 | 18.7 |
| | 2000 | 518 | 12.1% | 63.1 | 65.6 | 59.9 | 71.3 | 43.6 | 39.7 | 47.5 | 5.4 | 18.6 |
| | 2001 | 568 | 12.9% | 68.9 | 70.1 | 64.3 | 75.9 | 46.7 | 42.7 | 50.7 | 6.1 | 16.5 |
| | 2002 | 509 | 11.1% | 61.4 | 61.8 | 56.3 | 67.2 | 40.7 | 37.1 | 44.4 | 5.1 | 19.6 |
| | 2003 | 535 | 11.3% | 64.2 | 63.3 | 57.9 | 68.7 | 41.5 | 37.9 | 45.2 | 5.2 | 19.4 |
| 2004 | 593 | 12.6% | 70.9 | 68.2 | 62.6 | 73.7 | 44.2 | 40.5 | 47.9 | 5.2 | 19.4 | |
| 2005 | 516 | 11.0% | 61.1 | 59.0 | 53.9 | 64.2 | 39.1 | 35.7 | 42.6 | 4.9 | 20.6 | |
| Female | 1993 | 319 | 7.5% | 38.1 | 34.4 | 30.5 | 38.4 | 24.1 | 21.2 | 26.9 | 3.2 | 30.8 |
| | 1994 | 311 | 7.5% | 36.9 | 32.6 | 28.8 | 36.3 | 22.6 | 19.9 | 25.4 | 3.0 | 33.2 |
| | 1995 | 331 | 7.9% | 39.2 | 34.2 | 30.3 | 38.0 | 23.3 | 20.6 | 26.1 | 3.0 | 33.0 |
| | 1996 | 309 | 7.0% | 36.3 | 32.3 | 28.5 | 36.0 | 22.2 | 19.5 | 24.8 | 2.9 | 34.6 |
| | 1997 | 325 | 7.5% | 38.0 | 32.1 | 28.5 | 35.8 | 22.1 | 19.5 | 24.8 | 2.9 | 34.4 |
| | 1998 | 330 | 7.5% | 38.4 | 32.0 | 28.4 | 35.6 | 21.7 | 19.1 | 24.3 | 2.8 | 35.6 |
| | 1999 | 358 | 8.1% | 41.6 | 35.2 | 31.4 | 39.0 | 23.7 | 21.0 | 26.4 | 2.9 | 34.0 |
| | 2000 | 343 | 7.7% | 39.8 | 34.5 | 30.7 | 38.3 | 23.7 | 21.0 | 26.4 | 3.1 | 32.0 |
| | 2001 | 320 | 7.4% | 37.0 | 31.2 | 27.6 | 34.8 | 21.2 | 18.7 | 23.8 | 2.6 | 38.6 |
| | 2002 | 366 | 8.0% | 42.2 | 34.4 | 30.7 | 38.1 | 23.0 | 20.4 | 25.5 | 3.0 | 32.9 |
| | 2003 | 359 | 7.6% | 41.3 | 34.2 | 30.5 | 37.9 | 23.5 | 20.9 | 26.2 | 2.8 | 35.9 |
| 2004 | 376 | 8.0% | 43.0 | 34.2 | 30.5 | 37.8 | 22.8 | 20.3 | 25.3 | 2.8 | 35.9 | |
| 2005 | 356 | 7.6% | 40.4 | 32.6 | 29.0 | 36.1 | 21.9 | 19.4 | 24.3 | 2.6 | 37.8 | |
| Male & Female | 1993 | 884 | 10.5% | 54.0 | 52.4 | 48.9 | 56.0 | 36.0 | 33.5 | 38.5 | 4.7 | 21.2 |
| | 1994 | 936 | 11.3% | 56.9 | 55.9 | 52.2 | 59.6 | 38.5 | 35.9 | 41.1 | 5.0 | 20.1 |
| | 1995 | 877 | 10.6% | 53.2 | 50.5 | 47.1 | 54.0 | 34.2 | 31.8 | 36.6 | 4.4 | 22.5 |
| | 1996 | 859 | 9.9% | 51.7 | 49.1 | 45.8 | 52.5 | 33.3 | 30.9 | 35.6 | 4.4 | 22.7 |
| | 1997 | 877 | 10.3% | 52.5 | 49.3 | 46.0 | 52.7 | 33.5 | 31.2 | 35.9 | 4.3 | 23.1 |
| | 1998 | 892 | 10.4% | 53.2 | 49.0 | 45.7 | 52.3 | 33.0 | 30.7 | 35.4 | 4.3 | 23.5 |
| | 1999 | 869 | 10.1% | 51.8 | 47.4 | 44.2 | 50.7 | 31.7 | 29.4 | 33.9 | 4.0 | 24.9 |
| | 2000 | 861 | 9.9% | 51.2 | 47.7 | 44.4 | 50.9 | 32.3 | 30.0 | 34.6 | 4.2 | 24.0 |
| | 2001 | 888 | 10.2% | 52.6 | 47.9 | 44.7 | 51.1 | 32.4 | 30.1 | 34.6 | 4.2 | 23.9 |
| | 2002 | 875 | 9.6% | 51.6 | 46.1 | 43.0 | 49.2 | 30.7 | 28.6 | 32.9 | 4.0 | 25.0 |
| | 2003 | 894 | 9.5% | 52.5 | 46.4 | 43.3 | 49.6 | 31.2 | 29.1 | 33.4 | 3.9 | 25.8 |
| 2004 | 969 | 10.3% | 56.7 | 48.8 | 45.7 | 52.0 | 32.2 | 30.0 | 34.4 | 3.9 | 25.7 | |
| 2005 | 872 | 9.3% | 50.6 | 44.1 | 41.1 | 47.1 | 29.5 | 27.4 | 31.6 | 3.7 | 27.2 | |

Incidence rates for lung cancer (C33,C34)

