

Number of cases and incidence rates by sex and year of diagnosis
Brain and other CNS (C70-C72)

Male

Year of diagnosis	Number of cases	Percentage of all cancers ex. NMSC	Crude rate per 100,000 males	European age standardised incidence rate per 100,000 males (95% CI)	World age standardised incidence rate per 100,000 males (95% CI)
1993	61	2.0	7.6	8.3 (6.2,10.4)	7.2 (5.3,9.0)
1994	56	1.8	7.0	7.6 (5.6,9.6)	6.1 (4.4,7.7)
1995	55	1.8	6.8	7.6 (5.6,9.6)	6.3 (4.6,8.0)
1996	74	2.3	9.1	9.7 (7.5,12.0)	8.1 (6.2,10.0)
1997	61	2.0	7.5	8.0 (6.0,10.1)	6.3 (4.7,7.9)
1998	65	2.1	7.9	8.6 (6.5,10.6)	6.8 (5.1,8.5)
1999	58	1.9	7.1	7.5 (5.5,9.4)	6.1 (4.5,7.7)
2000	63	2.0	7.7	7.9 (5.9,9.9)	6.7 (5.0,8.4)
2001	72	2.2	8.7	8.9 (6.8,11.0)	6.8 (5.2,8.5)
2002	78	2.4	9.4	9.5 (7.4,11.6)	7.8 (6.0,9.7)
2003	70	2.0	8.4	8.5 (6.5,10.6)	7.1 (5.3,8.8)
2004	50	1.4	6.0	6.0 (4.3,7.7)	4.9 (3.5,6.3)
2005	77	2.1	9.1	8.9 (6.9,10.9)	7.2 (5.5,9.0)
2006	78	2.1	9.1	9.2 (7.1,11.2)	7.4 (5.7,9.1)
2007	81	2.0	9.4	9.3 (7.3,11.3)	7.9 (6.0,9.7)
2008	76	1.9	8.7	8.6 (6.6,10.5)	7.2 (5.4,8.9)

Female

Year of diagnosis	Number of cases	Percentage of all cancers ex. NMSC	Crude rate per 100,000 females	European age standardised incidence rate per 100,000 females (95% CI)	World age standardised incidence rate per 100,000 females (95% CI)
1993	44	1.4	5.3	5.4 (3.8,7.1)	5.0 (3.4,6.5)
1994	61	2.0	7.2	7.4 (5.5,9.3)	6.5 (4.8,8.2)
1995	25	0.8	3.0	2.7 (1.6,3.8)	2.2 (1.2,3.2)
1996	47	1.4	5.5	5.3 (3.8,6.9)	4.8 (3.3,6.4)
1997	53	1.6	6.2	6.0 (4.4,7.7)	5.0 (3.6,6.4)
1998	66	2.0	7.7	7.3 (5.5,9.2)	6.0 (4.4,7.6)
1999	31	0.9	3.6	3.6 (2.3,4.8)	3.3 (2.0,4.5)
2000	50	1.5	5.8	5.3 (3.8,6.9)	4.1 (2.9,5.4)
2001	56	1.7	6.5	6.2 (4.5,7.9)	5.1 (3.7,6.6)
2002	52	1.5	6.0	5.6 (4.0,7.1)	4.8 (3.3,6.2)
2003	54	1.5	6.2	6.2 (4.5,7.8)	5.5 (3.9,7.1)
2004	48	1.3	5.5	5.3 (3.7,6.8)	4.7 (3.2,6.1)
2005	44	1.2	5.0	4.6 (3.2,6.0)	3.8 (2.6,5.0)
2006	58	1.6	6.5	6.2 (4.6,7.9)	5.2 (3.7,6.6)
2007	54	1.4	6.0	5.6 (4.1,7.1)	5.0 (3.5,6.5)
2008	49	1.2	5.4	4.7 (3.4,6.1)	3.7 (2.5,4.9)

All persons

Year of diagnosis	Number of cases	Percentage of all cancers ex. NMSC	Crude rate per 100,000 persons	European age standardised incidence rate per 100,000 persons (95% CI)	World age standardised incidence rate per 100,000 persons (95% CI)
1993	105	1.7	6.4	6.8 (5.4,8.1)	6.0 (4.8,7.2)
1994	117	1.9	7.1	7.4 (6.0,8.8)	6.2 (5.0,7.4)
1995	80	1.3	4.9	5.1 (3.9,6.2)	4.2 (3.2,5.2)
1996	121	1.9	7.3	7.4 (6.1,8.8)	6.4 (5.2,7.6)
1997	114	1.8	6.8	6.9 (5.6,8.2)	5.6 (4.5,6.7)
1998	131	2.0	7.8	8.0 (6.6,9.4)	6.4 (5.3,7.6)
1999	89	1.4	5.3	5.4 (4.2,6.5)	4.6 (3.6,5.6)
2000	113	1.7	6.7	6.7 (5.4,7.9)	5.4 (4.4,6.5)
2001	128	1.9	7.6	7.4 (6.1,8.7)	5.9 (4.8,7.0)
2002	130	1.9	7.7	7.4 (6.1,8.7)	6.2 (5.1,7.4)
2003	124	1.8	7.3	7.2 (5.9,8.4)	6.2 (5.0,7.4)
2004	98	1.4	5.7	5.6 (4.5,6.8)	4.8 (3.8,5.8)
2005	121	1.7	7.0	6.6 (5.4,7.8)	5.5 (4.4,6.5)
2006	136	1.8	7.8	7.6 (6.3,8.9)	6.2 (5.1,7.4)
2007	135	1.7	7.7	7.4 (6.1,8.6)	6.4 (5.2,7.6)
2008	125	1.5	7.0	6.6 (5.4,7.8)	5.4 (4.4,6.4)

NOTES:

NMSC: Non-melanoma skin cancer

CI: Confidence interval

CNS: Central nervous system

Brain and other CNS (C70-C72)
(Trends in number of cases per year)



Brain and other CNS (C70-C72)
(Trends in age-standardised incidence rates)

