

Number of cases and incidence rates by sex and year of diagnosis
Breast (C50)

Male

Year of diagnosis	Number of cases	Percentage of all cancers ex.		Crude rate per 100,000 males	European age standardised incidence rate per 100,000 males (95% CI)	World age standardised incidence rate per 100,000 males (95% CI)
		NMSC				
1993	6	0.2		0.8	0.9 (0.2,1.6)	0.6 (0.1,1.1)
1994	6	0.2		0.7	0.9 (0.2,1.6)	0.5 (0.1,0.9)
1995	10	0.3		1.2	1.3 (0.5,2.1)	0.8 (0.3,1.3)
1996	8	0.3		1.0	1.0 (0.3,1.8)	0.7 (0.2,1.1)
1997	2	0.1		0.2	0.3 (0.0,0.6)	0.2 (0.0,0.5)
1998	7	0.2		0.9	0.9 (0.2,1.6)	0.5 (0.1,0.9)
1999	8	0.3		1.0	1.0 (0.3,1.7)	0.7 (0.2,1.2)
2000	6	0.2		0.7	0.7 (0.1,1.3)	0.5 (0.1,0.9)
2001	8	0.2		1.0	1.0 (0.3,1.7)	0.7 (0.2,1.1)
2002	9	0.3		1.1	1.1 (0.4,1.8)	0.7 (0.2,1.2)
2003	7	0.2		0.8	0.8 (0.2,1.4)	0.5 (0.1,0.9)
2004	2	0.1		0.2	0.2 (0.0,0.6)	0.1 (0.0,0.3)
2005	6	0.2		0.7	0.7 (0.1,1.3)	0.5 (0.1,0.9)
2006	2	0.1		0.2	0.2 (0.0,0.4)	0.1 (0.0,0.2)
2007	1	0.0		0.1	0.1 (0.0,0.3)	0.1 (0.0,0.3)
2008	8	0.2		0.9	0.8 (0.2,1.4)	0.5 (0.1,0.9)

Female

Year of diagnosis	Number of cases	Percentage of all cancers ex.		Crude rate per 100,000 females	European age standardised incidence rate per 100,000 females (95% CI)	World age standardised incidence rate per 100,000 females (95% CI)
		NMSC				
1993	771	24.2		92.1	93.2 (86.3,100.0)	68.0 (62.8,73.1)
1994	830	26.7		98.6	100.5 (93.3,107.6)	73.0 (67.7,78.4)
1995	880	27.8		104.1	104.3 (97.2,111.5)	75.6 (70.2,80.9)
1996	849	26.0		99.7	99.7 (92.8,106.7)	73.1 (67.9,78.4)
1997	857	26.1		100.1	99.6 (92.7,106.5)	72.6 (67.4,77.8)
1998	908	27.4		105.7	104.4 (97.4,111.5)	76.5 (71.2,81.8)
1999	940	28.2		109.2	105.0 (98.0,112.0)	75.8 (70.6,81.0)
2000	950	28.0		110.1	106.2 (99.2,113.2)	77.6 (72.4,82.9)
2001	926	28.1		107.1	103.8 (96.9,110.7)	75.7 (70.6,80.9)
2002	952	27.3		109.7	102.5 (95.8,109.3)	75.1 (70.0,80.2)
2003	1,033	28.9		118.8	112.0 (105.0,119.1)	81.8 (76.6,87.1)
2004	1,136	31.1		130.0	119.0 (111.8,126.1)	86.9 (81.5,92.3)
2005	1,085	29.7		123.3	110.9 (104.0,117.7)	80.7 (75.5,85.8)
2006	986	26.6		111.0	99.7 (93.3,106.1)	72.7 (67.9,77.5)
2007	1,159	29.7		129.2	116.8 (109.9,123.8)	84.7 (79.5,89.9)
2008	1,156	29.0		127.9	116.6 (109.7,123.5)	85.5 (80.3,90.7)

All persons

Year of diagnosis	Number of cases	Percentage of all cancers ex.		Crude rate per 100,000 persons	European age standardised incidence rate per 100,000 persons (95% CI)	World age standardised incidence rate per 100,000 persons (95% CI)
		NMSC				
1993	777	12.3		47.5	50.5 (46.9,54.2)	36.5 (33.8,39.2)
1994	836	13.4		50.9	54.3 (50.5,58.0)	39.0 (36.2,41.8)
1995	890	14.3		54.0	56.7 (52.8,60.5)	40.6 (37.8,43.4)
1996	857	13.3		51.6	53.8 (50.1,57.5)	39.0 (36.2,41.7)
1997	859	13.6		51.4	53.4 (49.8,57.1)	38.5 (35.8,41.2)
1998	915	14.2		54.5	56.0 (52.3,59.7)	40.5 (37.8,43.3)
1999	948	14.9		56.5	56.6 (53.0,60.3)	40.4 (37.7,43.2)
2000	956	14.7		56.8	57.0 (53.3,60.7)	41.2 (38.4,43.9)
2001	934	14.2		55.3	55.5 (51.9,59.1)	40.1 (37.4,42.7)
2002	961	14.1		56.6	54.9 (51.4,58.5)	39.8 (37.2,42.4)
2003	1,040	14.7		61.1	59.6 (55.9,63.3)	43.1 (40.4,45.9)
2004	1,138	15.8		66.5	63.2 (59.5,66.9)	45.7 (42.9,48.5)
2005	1,091	15.0		63.3	59.1 (55.6,62.7)	42.6 (39.9,45.2)
2006	988	13.2		56.7	52.8 (49.4,56.1)	38.1 (35.6,40.6)
2007	1,160	14.6		65.9	61.7 (58.0,65.3)	44.3 (41.7,47.0)
2008	1,164	14.4		65.6	61.5 (57.9,65.0)	44.7 (42.0,47.4)


NOTES:

NMSC: Non-melanoma skin cancer

CI: Confidence interval

Breast (C50)

(Trends in number of cases per year)



Breast (C50)

(Trends in age-standardised incidence rates)

