

**Number of cases and incidence rates by sex and year of diagnosis
Hodgkin's lymphoma (C81)**

Male					
Year of diagnosis	Number of cases	Percentage of all cancers ex.		European age standardised incidence rate per 100,000 males (95% CI)	World age standardised incidence rate per 100,000 males (95% CI)
		NMSC	Crude rate per 100,000 males		
1993	31	1.0	3.9	4.0 (2.6,5.4)	3.5 (2.3,4.8)
1994	22	0.7	2.7	3.0 (1.7,4.2)	2.4 (1.4,3.5)
1995	28	0.9	3.5	3.7 (2.3,5.1)	3.1 (1.9,4.2)
1996	19	0.6	2.3	2.4 (1.3,3.5)	2.1 (1.1,3.0)
1997	20	0.7	2.5	2.6 (1.5,3.8)	2.2 (1.2,3.1)
1998	11	0.4	1.3	1.4 (0.5,2.2)	1.2 (0.5,2.0)
1999	27	0.9	3.3	3.2 (2.0,4.4)	2.9 (1.8,4.0)
2000	25	0.8	3.0	3.1 (1.9,4.3)	2.6 (1.6,3.7)
2001	23	0.7	2.8	2.7 (1.6,3.8)	2.6 (1.5,3.7)
2002	10	0.3	1.2	1.2 (0.5,2.0)	1.2 (0.4,2.0)
2003	19	0.5	2.3	2.2 (1.2,3.2)	2.0 (1.1,2.9)
2004	23	0.6	2.7	2.7 (1.6,3.9)	2.5 (1.4,3.6)
2005	33	0.9	3.9	3.8 (2.5,5.1)	3.2 (2.1,4.3)
2006	22	0.6	2.6	2.5 (1.5,3.5)	2.4 (1.4,3.4)
2007	30	0.7	3.5	3.5 (2.3,4.8)	3.1 (1.9,4.2)
2008	27	0.7	3.1	3.0 (1.9,4.2)	2.6 (1.6,3.7)

Female					
Year of diagnosis	Number of cases	Percentage of all cancers ex.		European age standardised incidence rate per 100,000 females (95% CI)	World age standardised incidence rate per 100,000 females (95% CI)
		NMSC	Crude rate per 100,000 females		
1993	18	0.6	2.1	2.1 (1.1,3.0)	2.0 (1.1,3.0)
1994	16	0.5	1.9	1.8 (0.9,2.7)	1.8 (0.9,2.7)
1995	16	0.5	1.9	1.9 (0.9,2.8)	1.7 (0.8,2.5)
1996	14	0.4	1.6	1.6 (0.8,2.5)	1.5 (0.7,2.3)
1997	11	0.3	1.3	1.3 (0.5,2.1)	1.3 (0.5,2.1)
1998	18	0.5	2.1	1.9 (1.0,2.9)	1.6 (0.8,2.4)
1999	14	0.4	1.6	1.6 (0.7,2.4)	1.5 (0.7,2.4)
2000	11	0.3	1.3	1.3 (0.5,2.1)	1.2 (0.5,1.9)
2001	17	0.5	2.0	2.0 (1.1,3.0)	1.8 (0.9,2.7)
2002	18	0.5	2.1	2.1 (1.1,3.0)	2.0 (1.0,3.0)
2003	19	0.5	2.2	2.1 (1.2,3.1)	2.2 (1.2,3.2)
2004	19	0.5	2.2	2.2 (1.2,3.1)	2.2 (1.2,3.2)
2005	17	0.5	1.9	1.9 (1.0,2.8)	1.7 (0.9,2.6)
2006	23	0.6	2.6	2.5 (1.5,3.5)	2.3 (1.3,3.3)
2007	33	0.8	3.7	3.4 (2.2,4.6)	3.1 (2.0,4.3)
2008	39	1.0	4.3	4.1 (2.8,5.4)	3.6 (2.4,4.8)

All persons					
Year of diagnosis	Number of cases	Percentage of all cancers ex.		European age standardised incidence rate per 100,000 persons (95% CI)	World age standardised incidence rate per 100,000 persons (95% CI)
		NMSC	Crude rate per 100,000 persons		
1993	49	0.8	3.0	3.0 (2.1,3.8)	2.7 (1.9,3.5)
1994	38	0.6	2.3	2.3 (1.6,3.0)	2.1 (1.4,2.7)
1995	44	0.7	2.7	2.7 (1.9,3.5)	2.3 (1.6,3.0)
1996	33	0.5	2.0	2.0 (1.3,2.6)	1.8 (1.2,2.4)
1997	31	0.5	1.9	1.9 (1.2,2.6)	1.7 (1.1,2.3)
1998	29	0.5	1.7	1.7 (1.1,2.3)	1.4 (0.9,2.0)
1999	41	0.6	2.4	2.4 (1.6,3.1)	2.2 (1.5,2.9)
2000	36	0.6	2.1	2.2 (1.5,2.9)	1.9 (1.2,2.5)
2001	40	0.6	2.4	2.4 (1.6,3.1)	2.2 (1.5,2.9)
2002	28	0.4	1.7	1.7 (1.0,2.3)	1.6 (1.0,2.2)
2003	38	0.5	2.2	2.1 (1.5,2.8)	2.1 (1.4,2.8)
2004	42	0.6	2.5	2.4 (1.7,3.2)	2.3 (1.6,3.0)
2005	50	0.7	2.9	2.8 (2.0,3.6)	2.4 (1.7,3.1)
2006	45	0.6	2.6	2.5 (1.8,3.3)	2.3 (1.6,3.1)
2007	63	0.8	3.6	3.5 (2.6,4.3)	3.1 (2.3,3.9)
2008	66	0.8	3.7	3.5 (2.7,4.4)	3.1 (2.3,3.9)

NOTES:

NMSC: Non-melanoma skin cancer

CI: Confidence interval

Hodgkin's lymphoma (C81)

(Trends in number of cases per year)



Hodgkin's lymphoma (C81)

(Trends in age-standardised incidence rates)

