

Number of cases and incidence rates by sex and year of diagnosis
Liver (C22)

Male

Year of diagnosis	Number of cases	Percentage of all cancers ex.		Crude rate per 100,000 males	European age standardised incidence rate per 100,000 males (95% CI)	World age standardised incidence rate per 100,000 males (95% CI)
		NMSC				
1993	32	1.0		4.0	4.5 (2.9,6.0)	2.9 (1.8,3.9)
1994	25	0.8		3.1	3.6 (2.2,5.0)	2.3 (1.4,3.3)
1995	32	1.0		4.0	4.4 (2.9,6.0)	3.1 (2.0,4.2)
1996	33	1.0		4.1	4.4 (2.9,5.9)	3.0 (1.9,4.1)
1997	32	1.1		3.9	4.0 (2.6,5.4)	2.7 (1.7,3.6)
1998	26	0.8		3.2	3.3 (2.0,4.6)	2.4 (1.4,3.3)
1999	19	0.6		2.3	2.6 (1.4,3.8)	2.0 (1.1,2.9)
2000	34	1.1		4.1	4.2 (2.8,5.6)	2.9 (1.9,3.9)
2001	17	0.5		2.1	2.2 (1.1,3.2)	1.4 (0.7,2.0)
2002	41	1.2		4.9	4.9 (3.4,6.5)	3.3 (2.2,4.3)
2003	34	1.0		4.1	4.1 (2.7,5.5)	2.9 (1.8,3.9)
2004	31	0.9		3.7	3.7 (2.4,5.0)	2.5 (1.6,3.4)
2005	44	1.2		5.2	4.9 (3.5,6.4)	3.3 (2.3,4.3)
2006	54	1.4		6.3	6.1 (4.4,7.7)	3.9 (2.8,5.0)
2007	56	1.4		6.5	6.3 (4.6,7.9)	4.4 (3.2,5.7)
2008	46	1.1		5.3	4.9 (3.5,6.4)	3.4 (2.4,4.4)

Female

Year of diagnosis	Number of cases	Percentage of all cancers ex.		Crude rate per 100,000 females	European age standardised incidence rate per 100,000 females (95% CI)	World age standardised incidence rate per 100,000 females (95% CI)
		NMSC				
1993	33	1.0		3.9	3.2 (2.0,4.3)	2.0 (1.3,2.8)
1994	29	0.9		3.4	2.7 (1.6,3.7)	1.7 (1.0,2.4)
1995	17	0.5		2.0	1.7 (0.9,2.6)	1.2 (0.6,1.9)
1996	30	0.9		3.5	2.6 (1.6,3.6)	1.7 (1.1,2.4)
1997	27	0.8		3.2	2.1 (1.2,3.0)	1.3 (0.7,1.9)
1998	20	0.6		2.3	2.0 (1.1,3.0)	1.4 (0.7,2.1)
1999	28	0.8		3.3	2.3 (1.4,3.2)	1.4 (0.8,2.0)
2000	30	0.9		3.5	2.9 (1.8,3.9)	2.0 (1.2,2.8)
2001	8	0.2		0.9	0.9 (0.3,1.5)	0.6 (0.2,1.1)
2002	26	0.7		3.0	2.5 (1.5,3.5)	1.8 (1.0,2.6)
2003	19	0.5		2.2	1.6 (0.9,2.4)	1.3 (0.5,2.0)
2004	17	0.5		1.9	1.5 (0.8,2.3)	1.0 (0.5,1.6)
2005	23	0.6		2.6	2.0 (1.1,2.8)	1.4 (0.8,2.0)
2006	33	0.9		3.7	2.6 (1.7,3.6)	1.7 (1.0,2.3)
2007	29	0.7		3.2	2.4 (1.5,3.4)	1.6 (1.0,2.3)
2008	23	0.6		2.5	1.9 (1.1,2.6)	1.3 (0.7,2.0)

All persons

Year of diagnosis	Number of cases	Percentage of all cancers ex.		Crude rate per 100,000 persons	European age standardised incidence rate per 100,000 persons (95% CI)	World age standardised incidence rate per 100,000 persons (95% CI)
		NMSC				
1993	65	1.0		4.0	3.7 (2.8,4.6)	2.4 (1.8,3.0)
1994	54	0.9		3.3	3.0 (2.2,3.9)	2.0 (1.4,2.6)
1995	49	0.8		3.0	2.9 (2.1,3.8)	2.1 (1.4,2.7)
1996	63	1.0		3.8	3.4 (2.5,4.3)	2.3 (1.7,2.9)
1997	59	0.9		3.5	3.0 (2.2,3.8)	2.0 (1.4,2.5)
1998	46	0.7		2.7	2.6 (1.8,3.4)	1.8 (1.3,2.4)
1999	47	0.7		2.8	2.5 (1.8,3.3)	1.7 (1.2,2.3)
2000	64	1.0		3.8	3.5 (2.6,4.4)	2.4 (1.8,3.0)
2001	25	0.4		1.5	1.4 (0.8,1.9)	0.9 (0.5,1.3)
2002	67	1.0		3.9	3.5 (2.7,4.4)	2.4 (1.8,3.1)
2003	53	0.8		3.1	2.8 (2.0,3.5)	2.0 (1.4,2.6)
2004	48	0.7		2.8	2.5 (1.8,3.2)	1.7 (1.2,2.2)
2005	67	0.9		3.9	3.3 (2.5,4.1)	2.2 (1.7,2.8)
2006	87	1.2		5.0	4.2 (3.3,5.1)	2.7 (2.1,3.3)
2007	85	1.1		4.8	4.2 (3.3,5.1)	2.9 (2.2,3.6)
2008	69	0.9		3.9	3.3 (2.5,4.1)	2.3 (1.7,2.9)

NOTES:

NMSC: Non-melanoma skin cancer

CI: Confidence interval

Liver (C22)
(Trends in number of cases per year)



Liver (C22)

(Trends in age-standardised incidence rates)

