

**Number of cases and incidence rates by sex and year of diagnosis**  
**Oesophagus (C15)**

**Male**

Year of diagnosis	Number of cases	Percentage of all cancers ex.		Crude rate per 100,000 males	European age standardised incidence rate per 100,000 males (95% CI)	World age standardised incidence rate per 100,000 males (95% CI)
		NMSC				
1993	88	2.8		11.0	12.2 (9.6,14.8)	8.3 (6.5,10.1)
1994	100	3.2		12.5	13.9 (11.2,16.7)	9.4 (7.5,11.3)
1995	96	3.1		11.9	13.5 (10.8,16.2)	9.6 (7.6,11.6)
1996	79	2.5		9.7	10.3 (8.0,12.6)	6.8 (5.3,8.4)
1997	97	3.2		11.9	13.3 (10.6,16.0)	9.2 (7.3,11.1)
1998	100	3.2		12.2	13.5 (10.8,16.1)	9.2 (7.4,11.1)
1999	92	3.1		11.2	12.1 (9.6,14.6)	8.2 (6.4,9.9)
2000	70	2.2		8.5	9.0 (6.9,11.1)	6.0 (4.6,7.5)
2001	102	3.1		12.4	13.0 (10.4,15.5)	8.7 (6.9,10.4)
2002	84	2.5		10.1	10.5 (8.3,12.8)	7.0 (5.5,8.6)
2003	94	2.7		11.3	11.3 (9.0,13.6)	7.6 (6.0,9.2)
2004	107	3.0		12.8	12.8 (10.3,15.2)	8.6 (6.9,10.3)
2005	111	3.1		13.1	13.0 (10.6,15.5)	8.9 (7.2,10.6)
2006	122	3.2		14.3	13.7 (11.2,16.1)	9.1 (7.4,10.8)
2007	127	3.2		14.7	14.2 (11.7,16.7)	9.6 (7.9,11.3)
2008	125	3.1		14.4	13.8 (11.4,16.3)	9.7 (7.9,11.4)

**Female**

Year of diagnosis	Number of cases	Percentage of all cancers ex.		Crude rate per 100,000 females	European age standardised incidence rate per 100,000 females (95% CI)	World age standardised incidence rate per 100,000 females (95% CI)
		NMSC				
1993	56	1.8		6.7	5.3 (3.8,6.7)	3.4 (2.4,4.4)
1994	52	1.7		6.2	4.6 (3.3,5.9)	2.9 (2.0,3.9)
1995	56	1.8		6.6	5.2 (3.7,6.6)	3.3 (2.3,4.3)
1996	60	1.8		7.0	5.3 (3.9,6.7)	3.3 (2.4,4.2)
1997	74	2.3		8.6	6.3 (4.8,7.8)	4.0 (2.9,5.0)
1998	53	1.6		6.2	4.4 (3.1,5.7)	2.8 (1.9,3.6)
1999	61	1.8		7.1	5.6 (4.1,7.1)	3.7 (2.7,4.7)
2000	69	2.0		8.0	6.2 (4.6,7.7)	4.0 (3.0,5.1)
2001	63	1.9		7.3	5.4 (4.0,6.8)	3.5 (2.5,4.5)
2002	58	1.7		6.7	4.9 (3.6,6.2)	3.1 (2.2,4.0)
2003	52	1.5		6.0	4.5 (3.2,5.8)	3.0 (2.1,3.9)
2004	53	1.5		6.1	4.1 (2.9,5.2)	2.5 (1.7,3.3)
2005	49	1.3		5.6	4.0 (2.8,5.2)	2.5 (1.7,3.3)
2006	58	1.6		6.5	5.2 (3.8,6.6)	3.4 (2.4,4.4)
2007	75	1.9		8.4	6.1 (4.6,7.5)	4.0 (3.0,5.0)
2008	61	1.5		6.7	4.6 (3.4,5.8)	2.9 (2.1,3.7)

**All persons**

Year of diagnosis	Number of cases	Percentage of all cancers ex.		Crude rate per 100,000 persons	European age standardised incidence rate per 100,000 persons (95% CI)	World age standardised incidence rate per 100,000 persons (95% CI)
		NMSC				
1993	144	2.3		8.8	8.3 (6.9,9.7)	5.6 (4.6,6.6)
1994	152	2.4		9.2	8.8 (7.3,10.2)	5.9 (4.9,6.9)
1995	152	2.4		9.2	9.1 (7.6,10.5)	6.3 (5.2,7.3)
1996	139	2.2		8.4	7.5 (6.2,8.8)	4.9 (4.0,5.8)
1997	171	2.7		10.2	9.6 (8.1,11.1)	6.4 (5.4,7.5)
1998	153	2.4		9.1	8.5 (7.2,9.9)	5.8 (4.8,6.7)
1999	153	2.4		9.1	8.5 (7.1,9.9)	5.8 (4.8,6.7)
2000	139	2.1		8.3	7.5 (6.2,8.8)	5.0 (4.1,5.9)
2001	165	2.5		9.8	8.8 (7.4,10.2)	5.9 (4.9,6.8)
2002	142	2.1		8.4	7.4 (6.1,8.6)	4.9 (4.0,5.7)
2003	146	2.1		8.6	7.5 (6.2,8.7)	5.1 (4.2,5.9)
2004	160	2.2		9.4	8.1 (6.8,9.4)	5.4 (4.5,6.3)
2005	160	2.2		9.3	8.2 (6.9,9.5)	5.5 (4.6,6.5)
2006	180	2.4		10.3	9.0 (7.6,10.3)	6.0 (5.1,6.9)
2007	202	2.5		11.5	9.7 (8.4,11.1)	6.6 (5.6,7.6)
2008	186	2.3		10.5	9.0 (7.7,10.3)	6.1 (5.2,7.1)

**NOTES:**

NMSC: Non-melanoma skin cancer

CI: Confidence interval

# Oesophagus (C15)

(Trends in number of cases per year)



# Oesophagus (C15)

(Trends in age-standardised incidence rates)

