

## Number of cases and incidence rates by sex and year of diagnosis

### Ovary (C56)

#### Female

| Year of diagnosis | Number of cases | Percentage of all cancers ex. NMSC | Crude rate per 100,000 females | European age standardised incidence rate per 100,000 females (95% CI) | World age standardised incidence rate per 100,000 females (95% CI) |
|-------------------|-----------------|------------------------------------|--------------------------------|---|--|
| 1993              | 151             | 4.7                                | 18.0                           | 17.9 (14.9,20.9)  | 13.2 (11.0,15.5)   |
| 1994              | 155             | 5.0                                | 18.4                           | 18.2 (15.2,21.2)  | 13.4 (11.2,15.7)   |
| 1995              | 156             | 4.9                                | 18.5                           | 18.3 (15.3,21.3)  | 13.5 (11.3,15.8)   |
| 1996              | 148             | 4.5                                | 17.4                           | 16.9 (14.1,19.7)  | 12.4 (10.2,14.5)   |
| 1997              | 181             | 5.5                                | 21.2                           | 20.7 (17.5,23.8)  | 15.3 (12.9,17.7)   |
| 1998              | 183             | 5.5                                | 21.3                           | 20.6 (17.5,23.7)  | 15.2 (12.9,17.6)   |
| 1999              | 178             | 5.3                                | 20.7                           | 20.9 (17.8,24.1)  | 15.9 (13.4,18.3)   |
| 2000              | 167             | 4.9                                | 19.4                           | 18.4 (15.5,21.3)  | 13.5 (11.3,15.7)   |
| 2001              | 179             | 5.4                                | 20.7                           | 19.3 (16.4,22.3)  | 14.7 (12.4,17.0)   |
| 2002              | 190             | 5.5                                | 21.9                           | 19.8 (16.9,22.8)  | 14.6 (12.4,16.9)   |
| 2003              | 198             | 5.5                                | 22.8                           | 20.3 (17.3,23.2)  | 14.7 (12.5,17.0)   |
| 2004              | 197             | 5.4                                | 22.5                           | 20.0 (17.1,22.9)  | 14.9 (12.7,17.1)   |
| 2005              | 187             | 5.1                                | 21.2                           | 18.9 (16.1,21.7)  | 13.8 (11.7,15.9)   |
| 2006              | 190             | 5.1                                | 21.4                           | 19.2 (16.4,22.0)  | 14.2 (12.1,16.4)   |
| 2007*             | 156             | 4.0                                | 17.4                           | 15.0 (12.6,17.5)  | 10.8 (8.9,12.6)  |
| 2008*             | 176             | 4.4                                | 19.5                           | 16.8 (14.2,19.3)  | 12.0 (10.1,13.9)   |

#### All persons

| Year of diagnosis | Number of cases | Percentage of all cancers ex. NMSC | Crude rate per 100,000 persons | European age standardised incidence rate per 100,000 persons (95% CI) | World age standardised incidence rate per 100,000 persons (95% CI) |
|-------------------|-----------------|------------------------------------|--------------------------------|---|--|
| 1993              | 151             | 2.4                                | 9.2                            | 9.7 (8.1,11.2)  | 7.0 (5.9,8.2)  |
| 1994              | 155             | 2.5                                | 9.4                            | 9.8 (8.2,11.4)  | 7.2 (6.0,8.4)  |
| 1995              | 156             | 2.5                                | 9.5                            | 9.8 (8.3,11.4)  | 7.2 (6.0,8.4)  |
| 1996              | 148             | 2.3                                | 8.9                            | 9.1 (7.6,10.6)  | 6.6 (5.5,7.7)  |
| 1997              | 181             | 2.9                                | 10.8                           | 11.1 (9.4,12.7)   | 8.1 (6.9,9.4)  |
| 1998              | 183             | 2.8                                | 10.9                           | 11.0 (9.4,12.6)   | 8.1 (6.8,9.3)  |
| 1999              | 178             | 2.8                                | 10.6                           | 11.0 (9.4,12.7)   | 8.3 (7.0,9.5)  |
| 2000              | 167             | 2.6                                | 9.9                            | 9.9 (8.3,11.4)  | 7.1 (6.0,8.3)  |
| 2001              | 179             | 2.7                                | 10.6                           | 10.3 (8.8,11.8)   | 7.7 (6.5,8.9)  |
| 2002              | 190             | 2.8                                | 11.2                           | 10.6 (9.1,12.2)   | 7.7 (6.6,8.9)  |
| 2003              | 198             | 2.8                                | 11.6                           | 10.9 (9.3,12.4)   | 7.8 (6.7,9.0)  |
| 2004              | 197             | 2.7                                | 11.5                           | 10.7 (9.1,12.2)   | 7.8 (6.7,9.0)  |
| 2005              | 187             | 2.6                                | 10.8                           | 10.1 (8.6,11.5)   | 7.3 (6.2,8.4)  |
| 2006              | 190             | 2.5                                | 10.9                           | 10.1 (8.7,11.6)   | 7.4 (6.3,8.6)  |
| 2007*             | 156             | 2.0                                | 8.9                            | 8.0 (6.7,9.3)   | 5.7 (4.7,6.6)  |
| 2008*             | 176             | 2.2                                | 9.9                            | 8.9 (7.6,10.3)  | 6.3 (5.3,7.3)  |

#### NOTES:


NMSC: Non-melanoma skin cancer

CI: Confidence interval

\* A change in classification occurred in 2007 in which some ovarian cancers previously considered malignant are now considered to be non-malignant. The reduction in numbers between 2006 and 2007 may in part be due to this change

# Ovary (C56)

(Trends in number of cases per year)



## Ovary (C56)

(Trends in age-standardised incidence rates)

