

**Number of cases and incidence rates by sex and year of diagnosis  
Pancreas (C25)**

<b>Male</b>						
<b>Year of diagnosis</b>	<b>Number of cases</b>	<b>Percentage of all cancers ex.</b>		<b>Crude rate per 100,000 males</b>	<b>European age standardised incidence rate per 100,000 males (95% CI)</b>	<b>World age standardised incidence rate per 100,000 males (95% CI)</b>
		<b>NMSC</b>	<b>NMSC</b>			
1993	90	2.9	11.3	12.1 (9.6,14.6)	8.1 (6.3,9.8)	
1994	62	2.0	7.7	8.5 (6.4,10.7)	5.7 (4.3,7.2)	
1995	57	1.9	7.1	7.5 (5.5,9.5)	4.8 (3.5,6.1)	
1996	92	2.9	11.4	11.9 (9.5,14.4)	7.6 (6.0,9.3)	
1997	80	2.6	9.8	10.5 (8.2,12.9)	6.9 (5.3,8.4)	
1998	71	2.3	8.7	9.3 (7.1,11.5)	6.3 (4.8,7.8)	
1999	63	2.1	7.7	8.2 (6.2,10.3)	5.3 (4.0,6.7)	
2000	64	2.0	7.8	8.2 (6.2,10.2)	5.3 (3.9,6.6)	
2001	96	2.9	11.6	12.0 (9.6,14.5)	8.0 (6.4,9.7)	
2002	83	2.5	10.0	9.8 (7.7,11.9)	6.4 (4.9,7.8)	
2003	60	1.7	7.2	7.0 (5.2,8.8)	4.6 (3.4,5.8)	
2004	75	2.1	9.0	8.8 (6.8,10.9)	6.0 (4.6,7.4)	
2005	100	2.8	11.8	11.8 (9.5,14.2)	7.9 (6.3,9.5)	
2006	104	2.8	12.2	11.8 (9.5,14.0)	8.0 (6.4,9.6)	
2007	107	2.7	12.4	11.9 (9.6,14.1)	7.9 (6.3,9.4)	
2008	111	2.7	12.7	11.6 (9.4,13.8)	7.5 (6.0,8.9)	

<b>Female</b>						
<b>Year of diagnosis</b>	<b>Number of cases</b>	<b>Percentage of all cancers ex.</b>		<b>Crude rate per 100,000 females</b>	<b>European age standardised incidence rate per 100,000 females (95% CI)</b>	<b>World age standardised incidence rate per 100,000 females (95% CI)</b>
		<b>NMSC</b>	<b>NMSC</b>			
1993	85	2.7	10.2	7.9 (6.1,9.7)	5.3 (4.0,6.5)	
1994	78	2.5	9.3	7.5 (5.8,9.3)	5.0 (3.8,6.3)	
1995	65	2.1	7.7	5.9 (4.4,7.5)	3.9 (2.8,5.0)	
1996	73	2.2	8.6	6.7 (5.1,8.3)	4.2 (3.1,5.4)	
1997	82	2.5	9.6	7.2 (5.5,8.8)	4.7 (3.5,5.8)	
1998	65	2.0	7.6	5.6 (4.2,7.0)	3.6 (2.6,4.7)	
1999	58	1.7	6.7	5.0 (3.6,6.4)	3.2 (2.2,4.1)	
2000	81	2.4	9.4	7.7 (6.0,9.5)	5.3 (4.0,6.5)	
2001	75	2.3	8.7	6.2 (4.7,7.7)	4.0 (3.0,5.0)	
2002	87	2.5	10.0	7.5 (5.9,9.2)	4.8 (3.7,6.0)	
2003	97	2.7	11.2	8.0 (6.3,9.7)	5.2 (4.0,6.3)	
2004	83	2.3	9.5	7.2 (5.5,8.8)	4.8 (3.6,5.9)	
2005	81	2.2	9.2	6.3 (4.9,7.8)	4.0 (3.0,4.9)	
2006	103	2.8	11.6	8.4 (6.7,10.1)	5.5 (4.3,6.6)	
2007	85	2.2	9.5	6.9 (5.4,8.5)	4.6 (3.5,5.7)	
2008	111	2.8	12.3	8.9 (7.2,10.7)	5.8 (4.6,7.0)	

<b>All persons</b>						
<b>Year of diagnosis</b>	<b>Number of cases</b>	<b>Percentage of all cancers ex.</b>		<b>Crude rate per 100,000 persons</b>	<b>European age standardised incidence rate per 100,000 persons (95% CI)</b>	<b>World age standardised incidence rate per 100,000 persons (95% CI)</b>
		<b>NMSC</b>	<b>NMSC</b>			
1993	175	2.8	10.7	9.7 (8.3,11.2)	6.5 (5.4,7.5)	
1994	140	2.3	8.5	7.9 (6.6,9.3)	5.3 (4.4,6.3)	
1995	122	2.0	7.4	6.6 (5.4,7.8)	4.2 (3.4,5.1)	
1996	165	2.6	9.9	8.9 (7.5,10.2)	5.7 (4.7,6.6)	
1997	162	2.6	9.7	8.6 (7.3,10.0)	5.6 (4.7,6.5)	
1998	136	2.1	8.1	7.5 (6.2,8.8)	5.0 (4.1,5.9)	
1999	121	1.9	7.2	6.3 (5.2,7.5)	4.1 (3.3,4.9)	
2000	145	2.2	8.6	7.8 (6.5,9.1)	5.2 (4.3,6.1)	
2001	171	2.6	10.1	8.8 (7.4,10.2)	5.8 (4.9,6.7)	
2002	170	2.5	10.0	8.5 (7.2,9.9)	5.5 (4.6,6.4)	
2003	157	2.2	9.2	7.7 (6.4,8.9)	5.0 (4.1,5.8)	
2004	158	2.2	9.2	8.0 (6.7,9.3)	5.3 (4.4,6.2)	
2005	181	2.5	10.5	8.8 (7.5,10.2)	5.8 (4.9,6.7)	
2006	207	2.8	11.9	10.1 (8.7,11.5)	6.7 (5.7,7.7)	
2007	192	2.4	10.9	9.2 (7.8,10.5)	6.1 (5.2,7.0)	
2008	222	2.8	12.5	10.1 (8.7,11.4)	6.5 (5.6,7.4)	

**NOTES:**

NMSC: Non-melanoma skin cancer

CI: Confidence interval

**Pancreas (C25)**  
(Trends in number of cases per year)



**Pancreas (C25)**  
(Trends in age-standardised incidence rates)

