

Number of cases and incidence rates by sex and year of diagnosis

Prostate (C61)

Male

Year of diagnosis	Number of cases	Percentage of all cancers ex. NMSC	Crude rate per 100,000 males	European age standardised incidence rate per 100,000 males (95% CI)	World age standardised incidence rate per 100,000 males (95% CI)
1993	473	15.2	59.3	63.9 (58.0,69.8)	39.0 (35.3,42.7)
1994	486	15.6	60.6	64.5 (58.7,70.3)	39.4 (35.8,43.1)
1995	513	16.8	63.8	67.4 (61.4,73.3)	40.2 (36.6,43.8)
1996	462	14.6	57.0	60.3 (54.7,65.9)	37.1 (33.6,40.6)
1997	488	16.1	59.8	62.6 (57.0,68.2)	37.7 (34.3,41.2)
1998	502	16.0	61.3	63.5 (57.9,69.1)	38.9 (35.3,42.4)
1999	500	16.6	61.1	63.1 (57.5,68.7)	39.5 (35.9,43.1)
2000	590	18.9	71.9	73.2 (67.2,79.1)	46.3 (42.4,50.2)
2001	570	17.4	69.1	70.3 (64.5,76.1)	44.7 (40.9,48.5)
2002	679	20.5	81.9	82.3 (76.0,88.5)	53.3 (49.2,57.5)
2003	790	22.7	94.9	93.8 (87.2,100.4)	60.8 (56.4,65.2)
2004	809	22.8	96.7	94.6 (88.1,101.2)	62.0 (57.6,66.4)
2005	821	22.8	97.2	93.7 (87.2,100.1)	61.2 (56.8,65.5)
2006	854	22.7	100.1	95.6 (89.2,102.1)	63.4 (59.0,67.7)
2007	941	23.4	109.2	103.5 (96.9,110.2)	69.2 (64.6,73.7)
2008	962	23.6	110.5	103.8 (97.2,110.4)	69.4 (64.9,73.9)

All persons

Year of diagnosis	Number of cases	Percentage of all cancers ex. NMSC	Crude rate per 100,000 persons	European age standardised incidence rate per 100,000 persons (95% CI)	World age standardised incidence rate per 100,000 persons (95% CI)
1993	473	7.5	28.9	24.9 (22.6,27.2)	15.7 (14.1,17.2)
1994	486	7.8	29.6	25.3 (23.0,27.6)	15.9 (14.4,17.4)
1995	513	8.2	31.1	26.1 (23.7,28.4)	16.0 (14.5,17.5)
1996	462	7.2	27.8	24.0 (21.8,26.3)	15.2 (13.7,16.7)
1997	488	7.7	29.2	24.4 (22.2,26.6)	15.1 (13.7,16.6)
1998	502	7.8	29.9	25.4 (23.1,27.7)	16.0 (14.5,17.5)
1999	500	7.9	29.8	25.8 (23.5,28.2)	16.6 (15.0,18.2)
2000	590	9.1	35.1	30.7 (28.1,33.2)	19.9 (18.1,21.6)
2001	570	8.7	33.7	29.3 (26.9,31.8)	19.1 (17.5,20.8)
2002	679	10.0	40.0	35.0 (32.3,37.7)	23.2 (21.4,25.1)
2003	790	11.2	46.4	40.4 (37.5,43.3)	26.7 (24.7,28.7)
2004	809	11.2	47.3	41.2 (38.3,44.1)	27.6 (25.5,29.6)
2005	821	11.3	47.6	41.0 (38.1,43.9)	27.3 (25.3,29.3)
2006	854	11.4	49.0	42.7 (39.8,45.6)	28.7 (26.7,30.8)
2007	941	11.9	53.5	46.6 (43.6,49.6)	31.6 (29.5,33.8)
2008	962	11.9	54.2	47.1 (44.0,50.1)	31.9 (29.8,34.1)


NOTES:

NMSC: Non-melanoma skin cancer

CI: Confidence interval

Prostate (C61)

(Trends in number of cases per year)



Prostate (C61)

(Trends in age-standardised incidence rates)

