

 COMFORT B Scores <u>can</u> be used to assess sedation & comfort in patients with complex needs. When scoring each category ask yourself 'what is normal for this child?'

Ask their parent to tell you their normal! A grimace could be their happy face.



• DO NOT COMFORT score patients who are on neuromuscular blocking agents. The score is dependent on the interpretation of behavioural cues which cannot be displayed if the patient is muscle relaxed.

Assess COMFORT B Scores a minimum of 6 hourly.

3-4 hourly really is the optimum for patient comfort while not overloading the bedside nurse with extra work.

 If your patients' COMFORT B score is not in their set target range you must do something about it!



- If you make a change to sedation/ analgesia you must reassess the COMFORT score one hour later.
- Continue to assess COMFORT B score until point of extubation even if all sedative agents have been discontinued.
 - Patients can safely extubate with a COMFORT B score of 12-17

and when you extubate...

 Following extubation patients should continue to have their COMFORT B score assessed until at least 12- 24 hours after all sedative & opioid agents have been discontinued.

Eg if the patient extubated at 10am and all sedative agents were stopped at 10am the patient should continue to have COMFORT B scores measured until at least 10pm if not 10am the following day.

If the patient has been on IV/enteral **sedation for more than 5 days** it would be recommended to continue assessing COMFORT B scores for a **minimum of 24hours after all sedative agents have been discontinued**.